


Glucocorticoid Induced Osteoporosis

Developed by the Lebanese National Task Force for Osteoporosis and Metabolic Bone Disorders*

Endorsed by:

Lebanese Society for Osteoporosis and Metabolic Bone Disorders, Lebanese Society of Endocrinology, Lebanese Society of Obstetrics and Gynecology, Lebanese Society of Rheumatology, Lebanese Orthopedics Society, Lebanese Society of Radiology, Lebanese Society of Family Medicine, Lebanese Society of Internal Medicine, Lebanese Society of General Practitioners.



¹ Osteoporosis Canada guidelines 2010.

² ACR 2010 guidelines.

³ IOF-ECTS 2012 guidelines.

⁴ Prednisone or prednisolone.

⁵ 10-year risk for MOF adjustment: dose <2.5 mg/day, decrease FRAX by 20% ; dose 2.5-7.5 mg/day: no adjustment; Dose ≥ 7.5 mg/day: increase FRAX by 15%.

⁶ FDA approved therapies for GIOP: alendronate, risedronate, zoledronic acid and teriparatide.

⁷ Teriparatide is indicated in high risk individuals. High risk individuals are defined as postmenopausal women and men ≥ 50 years with high FRAX estimate as defined by FRAX Lebanon treatment thresholds, or premenopausal women and men < 50 years who have a history of fragility fracture and on a prednisone dose ≥7.5 mg daily for more than 3 months.

⁸ Treat for the duration of steroid therapy.

* Lebanese National Task Force for Osteoporosis and Metabolic Bone Disorders Members

Ghada El-Hajj Fuleihan, MD, MPH; Rafic Baddoura, MD, MPH; Marlene Chakhtoura, MD, MSc; Asma Arabi, MD, MSc; George Halaby, MD; Imad Uthman, MD; Jad Okais, MD;


Ibrahim Salti, MD, PhD; Muhieddine Seoud, MD; Asaad Taha, MD; Naji Attallah, MD.

Glucocorticoid Induced Osteoporosis

Developed by the Lebanese National Task Force for Osteoporosis and Metabolic Bone Disorders*

Endorsed by:

Lebanese Society for Osteoporosis and Metabolic Bone Disorders, Lebanese Society of Endocrinology, Lebanese Society of Obstetrics and Gynecology, Lebanese Society of Rheumatology, Lebanese Orthopedics Society, Lebanese Society of Radiology, Lebanese Society of Family Medicine, Lebanese Society of Internal Medicine, Lebanese Society of General Practitioners.



1 American College of Rheumatology (ACR) 2010 guidelines

2 Treat for the duration of steroid therapy.

3 If no previous fragility fracture, no recommendation was made by ACR. In a young individual with low bone mineral density and significant bone loss and who requires long term systemic glucocorticoid therapy, osteoporosis therapy may be suggested; expert opinion, Hansen et al, JBMR 2011.

* Lebanese National Task Force for Osteoporosis and Metabolic Bone Disorders Members

Ghada El-Hajj Fuleihan, MD, MPH; Rafic Baddoura, MD, MPH; Marlene Chakhtoura, MD, MSc; Asma Arabi, MD, MSc; George Halaby, MD; Imad Uthman, MD; Jad Okais, MD;

Ibrahim Salti, MD, PhD; Muhieddine Seoud, MD; Asaad Taha, MD; Najj Attallah, MD.