

ANESTHESIA FOR GINGIVECTOMY AND DENTAL EXTRACTIONS IN A CHILD WITH I-CELL DISEASE

- A Case Report -

ABDUL KADER M. MAHFOUZ* AND G. GEORGE**

Abstract

I-cell disease (mucopolipidosis II) is a rare metabolic disorder resulting from the deficiency of a specific lysosomal enzyme, N-acetylglucosamine-1-phosphotransferase. Developmental delay and growth failure are common presentations of I-cell disease. Psychomotor deterioration is rapid and progressive. Some physical signs, such as hip dislocations, inguinal hernia, hepatomegaly, joint limitation, and skin changes may be present at birth. Coarse facial features and skeletal abnormalities become more conspicuous with time. The life expectancy of children with this condition is poor, with death usually occurring around the fifth year. A case report of the anesthetic management of gingivectomy with multiple dental extractions in a 5 years old Omani female, with I-cell disease, is presented. The problems faced and their management during anesthesia are described.

Introduction

I-cell disease (Mucopolipidosis II) was first described by Leroy and DeMars in 1967¹. It is an autosomal recessive, metabolic storage disorder due to a deficiency of the enzyme N-acetylglucosamine-1-phosphotransferase, which is one of two enzymes involved in the biosynthesis of mannose-6-phosphate². This compound is a common marker enabling lysosomal enzymes to be transported to the lysosomal compartment of all cells. Without the transferase, lysosomal enzymes escape into the extra-cellular fluid resulting in marked elevation of lysosomal enzymes in plasma³. This transferase deficiency, and consequent lysosomal depletion, results in the accumulation of a number of macromolecules, mucopolysaccharides and mucolipids, in the lysosome. This can be seen as coarse cytoplasmic granular inclusions in cultured skin fibroblasts giving rise to the name 'I-cell' disease⁴.

The most common features of the condition are mental and physical retardation with typical orofacial features. Infants with I-cell disease are typically underweight at birth, with muscle hypotonia, and coarse facial features, the full clinical picture of the disorder presenting at between 6 and 8 months⁵. Other features include joint stiffness, dislocated hips and tight, thickened skin,

* MD, PhD, Sr. Consultant Anaesthesiologist.

** MD, DA, Jr. Specialist Anaesthesiologist.

From Anaesthesia Department, Al Nahdha Hospital, Sultanate of Oman.

Corresponding author: Dr. Abdul Kader M. Mahfouz, Senior Consultant Anaesthesiologist, Al Nahdha Hospital, Muscat, Sultanate of Oman. P.O. Box: 937, PC 112 Ruwi, GSM: +968-99360867, E mail: abdulkaderm2000@hotmail.com

umbilical and or inguinal hernia, hepatomegaly, aortic insufficiency, and hazy corneas⁶. The typical orofacial features in I-cell disease include: coarse facial features, high, narrow forehead, puffy eyelids, epicanthal folds, flat nasal bridge, anteverted nares, long philtrum, prominent gingival hyperplasia and macroglossia. Due to marked enlargement of the gingivae and alveolar process, the lower part of the face usually has a 'fish-like' profile⁷. The gingival enlargement is progressive, from 4 months of age, to the extent that when jaw closure is attempted, contact is only made in the posterior part of the mouth, giving rise to an apparently open bite.

Diagnosis of the condition is often made in retrospect as a result of physical and mental delay. However, the presence of marked elevations of lysosomal enzymes in the plasma is an accurate diagnostic test for this disorder together with the presence of large lysosomal inclusions in peripheral lymphocytes³.

The life expectancy of children with I-cell disease is poor, as death occurs between the fifth and seventh year from recurrent upper respiratory tract infections, bronchopneumonia and heart failure⁸. For children with I-cell disease, the only therapeutic approach currently available is bone marrow transplantation to supply a source of structurally normal lysosomal enzymes⁹.

The anesthetic consideration in such cases included a high risk of difficult intubation and possibility of requirements of postoperative mechanical ventilation after the end of surgical procedure.

Case report

An Omani female, 5 years and 11 months of age, was referred to Al Nahdha Hospital (tertiary Hospital for Oro-Maxillo-Facial services in Oman) from the Genetic Clinic where she was being followed-up regularly as a case of I-Cell Disease (Mucopolidosis Type II). She was the second of three siblings. One of them was well and free of the disease while the other child died at the age of 1½ years due to the same disease.

She had the history of chronic nasal discharge and noisy breathing and also used to get recurrent chest infections for which she was admitted to hospital twice

and treated with antibiotics and nebulisation.

On this occasion she presented with generalised gingival hyperplasia, poor oral hygiene and retained roots of lower anterior teeth. She was admitted for gingivectomy with multiple dental extractions under general anesthesia.

On examination she was found to have: coarse facies, a depressed nose, a prominent forehead, high arched palate, widened wrists, a very large tongue, carious teeth, hypertrophied gums, generalized hypotonia, umbilical hernia, hepatomegaly and thick skin. She weighed 7 kg and was noticed to have a developmental delay. Examination of the heart and lungs were unremarkable except for some conducted sounds over the lungs.

Anesthesia Management

The child was premedicated with midazolam 3.5 mg orally, and atropine 150 µg orally half an hour before the procedure. Due to the history of recurrent chest infection and presence of conducted sounds over the chest, the child was nebulized with salbutamol solution 1 mg and ipratropium solution 75 µg half an hour before the procedure as a prophylactic measure against bronchospasm during anesthesia. The child was kept fasting for 6 hours before the procedure with a continuous IV infusion of dextrose 4.3% in 0.18 % normal saline at the rate of 28 ml.hr⁻¹.

On arrival in the operating theatre, routine monitoring was commenced which included ECG, non-invasive blood pressure (NIBP), pulse oximetry (S_{P_O2}) and end-tidal CO₂ tension (ET_{CO₂}).

General anesthesia was induced with sevoflurane 6% in oxygen. The patient's airway became progressively obstructed in spite of the use of an oropharyngeal airway size 1 and the jaw thrust manoeuvre. It was also not possible to assist her respiration with a bag and mask. A size 2 LMA was introduced and this successfully maintained the airway. Anesthesia was deepened subsequently. Due to the risk of bleeding from the planned surgery, endotracheal intubation was deemed necessary to guard against the risk of aspiration. After an adequate depth of anesthesia was reached, laryngoscopy and intubation was attempted after removing the LMA. Laryngoscopy was very

difficult due to the stiffness of neck and it was not possible to properly position the patient for intubation. Because of this as well as the large tongue, the glottis could not be visualised. The LMA was reintroduced and another attempt of intubation was carried out with a Miller blade instead of the Macintosh blade, but it was still not possible to visualize the glottis. Once again the LMA was reinserted to maintain the airway. After instillation of vasoconstrictor nasal drops, intubation was attempted using a paediatric (3 mm) fibre-optic bronchoscope with a 4.5 mm uncuffed tracheal tube mounted on it. This had to be abandoned due to nasal bleeding from congested hypertrophied tissues and large adenoids, as well as rapid desaturation after removal of the LMA.

Finally, a sand bag was placed under the patient's shoulder to get some head extension and then the tip of epiglottis could be visualized. Oral intubation finally succeeded with a 4mm ID endotracheal tube using a stylet, pushing the endotracheal tube behind the barely visualised epiglottis. Correct positioning of the tube was confirmed by auscultation of chest and by capnography. The tube was secured by adhesive tape and a throat pack was inserted.

Anesthesia was maintained with sevoflurane in oxygen and nitrous oxide, with spontaneous respiration using a modified Jackson-Rees circuit. The respiration was assisted manually as required. At the end of the surgical procedure, the throat pack was removed and the pharynx cleared of secretions by suction. The patient was extubated when fully conscious and the recovery was uneventful.

Discussion

Anesthetising a patient with I-cell disease requires skill and should only be undertaken by an experienced anesthetist. Many factors contribute to difficult intubation including large tongue, limited

cervical movement and hypertrophy of nasal tissues, tonsils and adenoids due to the presence of inclusion bodies. As the children grow older the airway becomes gradually more restricted as the initially swollen passages become stiff and easily traumatized¹⁰. Because of the poor growth potential, atrophied muscles, hypotonia, poor compliance of the narrow thoracic cage and recurrent chest infection, the baseline respiratory status of these children is compromised and this may explain the need of postoperative mechanical ventilation in some cases¹¹.

As general anesthesia is considered risky in this disease, it is recommended that elective surgical procedures be avoided as far as possible. If a procedure is considered essential, it should be undertaken at a major medical facility where pediatric anesthesia and pediatric critical care services are available. There should also be facilities to perform fibre-optic intubation. A detailed explanation of the anesthetic risk should be given to the parents¹².

In the present case, the inability to maintain the airway without the use of a laryngeal mask airway limited the usefulness of the fibre-optic bronchoscope as a tool of intubation. In addition, bleeding occurred during a trial of fibre-optic intubation because of hypertrophied adenoids and congested nasal passage.

Although it was possible to maintain airway patency with a LMA, it was considered unsafe to perform the surgical procedure (gingivectomy and dental extraction) with a LMA in situ due to the risk of aspiration. FastTrack intubation using an intubating LMA was not possible in this case due to the unavailability of a suitable sized FastTrack LMA which matching the body weight of the child.

Many reports¹⁰⁻¹¹ have mentioned the possibility of difficult intubation as a risk factor during anesthesia, but in this case the inability to maintain the airway was an additional risk.

References

1. LEROY JG, DEMARS RI: Mutant enzymatic and cytological phenotypes in cultured human fibroblasts. *Science*; 1967, 157:804-6.
2. DIERKS T, SCHLOTAWA L, FRESE MA, RADHAKRISHNAN K, VON FIGURA K, SCHMIDT B: Molecular basis of multiple sulfatase deficiency, mucopolipidosis II/III and Niemann-Pick C1 disease-lysosomal storage disorders caused by defects of non-lysosomal proteins. *Biochimica et Biophysica Acta*; 2009, 1793(4):710-25.
3. CAREY WF, JAUZEMS A, RICHARDSON M, FONG BA, CHIN SJ, NELSON PV: Prenatal diagnosis of mucopolipidosis-electron microscopy and biochemical evaluation. *Prenatal Diagnosis*; 1999, 252-6.
4. HANAI J, LEORY J, O'BRIEN JS: Ultra structure of cultured fibroblasts in I-cell disease. *American Journal of Disabled Children*; 1971, 122:34-8.
5. SPITZ RA, DOUGHTY RA, SPACKMAN TJ, MURANE MJ, COATES PM, KOLDOVSKY O, ZACKAI EH: Neonatal presentation of I-cell disease. *Journal of Paediatrics*; 1978, 93:954-8.
6. GOODMAN RM, GORLIN RJ: Malformed infant and child: an illustrated guide. Oxford: Oxford University Press, 1983, 182-3.
7. GORLIN RJ, COHN MM, LEVIN SS: Syndromes of the Head and Neck, 3rd ed. Oxford: Oxford University Press, 1990, 124-6.
8. SHEIKH S, MADIRAJU K, QAZI Q, RAO M: Improved morbidity with the use of nasal continuous positive airway pressure in I-cell disease. *Pediatric pulmonology*; 1998, 25:128-9.
9. DUMONTEL C, CONSTANT H, SOUILLET G, ZABOT MT: In vitro toxic effects of certain antibodies on the fibroblasts of two children with I-cell disease. *Cell Biology and Toxicology*; 1998, 14:333-43.
10. BAINS DB, STREET N, OVERTON JH: Anaesthetic implications of mucopolipidosis. *Paediatric Anaesth*; 1993, 3:303:6.
11. BAUM VC, O'FLAHERTY JE: Anesthetic consideration in MLII. In: Anesthesia for genetic, metabolic and dysmorphic syndromes of childhood. 2nd ed. Lippincott Williams & Wilkins, 2006, 187.
12. KEMPTHORNE PM, BROWN TC: Anaesthesia and the mucopolysaccharidosis. A survey of techniques and problems. *Anaesth Intensive Care*; 1983, 11:203-7.